


*Statistics for Managers Using
Microsoft Excel*
7th Edition



Chapter 8

Confidence Interval Estimation



Learning Objectives

In this chapter, you learn:

- To construct and interpret confidence interval estimates for the mean and the proportion
- How to determine the sample size necessary to develop a confidence interval for the mean or proportion



Chapter Outline


Content of this chapter

- Confidence Intervals for the **Population Mean, μ**
 - when Population Standard Deviation σ is **Known**
 - when Population Standard Deviation σ is **Unknown**
- Confidence Intervals for the **Population Proportion, π**
- Determining the **Required Sample Size**

Point and Interval Estimates

DCOVA A

- A **point estimate** is a single number,
- a **confidence interval** provides additional information about the variability of the estimate



Point Estimates

We can estimate a Population Parameter ...		with a Sample Statistic (a Point Estimate)
Mean	μ	\bar{X}
Proportion	π	p



Confidence Intervals

DCOVA A

- How much uncertainty is associated with a point estimate of a population parameter?
- An **interval estimate** provides more information about a population characteristic than does a **point estimate**
- Such interval estimates are called **confidence intervals**



Confidence Interval Estimate

DCOVA

- An interval gives a **range** of values:
 - Takes into consideration variation in sample statistics from sample to sample
 - Based on observations from 1 sample
 - Gives information about closeness to unknown population parameters
 - Stated in terms of level of confidence
 - e.g. 95% confident, 99% confident
 - Can never be 100% confident



Confidence Interval Example

Cereal fill example

- Population has $\mu = 368$ and $\sigma = 15$.
- If you take a sample of size $n = 25$ you know
 - $368 \pm 1.96 * 15 / \sqrt{25} = (362.12, 373.88)$ contains 95% of the sample means
 - When you don't know μ , you use \bar{X} to estimate μ
 - If $\bar{X} = 362.3$ the interval is $362.3 \pm 1.96 * 15 / \sqrt{25} = (356.42, 368.18)$
 - Since $356.42 \leq \mu \leq 368.18$ the interval based on this sample makes a correct statement about μ .

But what about the intervals from other possible samples of size 25?

Confidence Interval Example

DCOVA

(continued)

Sample #	\bar{X}	Lower Limit	Upper Limit	Contain μ ?
1	362.30	356.42	368.18	Yes
2	369.50	363.62	375.38	Yes
3	360.00	354.12	365.88	No
4	362.12	356.24	368.00	Yes
5	373.88	368.00	379.76	Yes

Confidence Interval Example


DCOVA

(continued)

- In practice you only take one sample of size n
- In practice you do not know μ so you do not know if the interval actually contains μ
- However you do know that 95% of the intervals formed in this manner will contain μ
- Thus, based on the one sample, you actually selected you can be 95% confident your interval will contain μ (this is a 95% **confidence interval**)

Note: 95% confidence is based on the fact that we used $Z = 1.96$.

Estimation Process





General Formula

DCOVA A

- The general formula for all confidence intervals is:

$$\text{Point Estimate} \pm (\text{Critical Value})(\text{Standard Error})$$

Where:

- **Point Estimate** is the sample statistic estimating the population parameter of interest
- **Critical Value** is a table value based on the sampling distribution of the point estimate and the desired confidence level
- **Standard Error** is the standard deviation of the point estimate



Confidence Level

DCOVA A

- Confidence Level
 - Confidence the interval will contain the unknown population parameter
 - A percentage (less than 100%)


Confidence Level, $(1-\alpha)$

DCOVA

(continued)

- Suppose confidence level = 95%
- Also written $(1 - \alpha) = 0.95$, (so $\alpha = 0.05$)
- A relative frequency interpretation:
 - 95% of all the confidence intervals that can be constructed will contain the unknown true parameter
- A specific interval either will contain or will not contain the true parameter
 - No probability involved in a specific interval

Confidence Intervals



Confidence Interval for μ (σ Known)

- Assumptions
 - Population standard deviation σ is known
 - Population is normally distributed
 - If population is not normal, use large sample
- Confidence interval estimate:

$$\bar{X} \pm Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

where \bar{X} is the point estimate

$Z_{\alpha/2}$ is the normal distribution critical value for a probability of $\alpha/2$ in each tail

σ/\sqrt{n} is the standard error


Finding the Critical Value, $Z_{\alpha/2}$

DCOVA

$$Z_{\alpha/2} = \pm 1.96$$

- Consider a 95% confidence interval:

$$1 - \alpha = 0.95 \text{ so } \alpha = 0.05$$





Common Levels of Confidence

DCOVA A


- Commonly used confidence levels are 90%, 95%, and 99%

Confidence Level	Confidence Coefficient, $1 - \alpha$	$Z_{\alpha/2}$ value
80%	0.80	1.28
90%	0.90	1.645
95%	0.95	1.96
98%	0.98	2.33
99%	0.99	2.58
99.8%	0.998	3.08
99.9%	0.999	3.27

Intervals and Level of Confidence

DCOVA A


Sampling Distribution of the Mean



Intervals extend from

$$\bar{X} - Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

to

$$\bar{X} + Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$



Confidence Intervals

(1- α)x100% of intervals constructed contain μ ;
 (α) x100% do not.

Example

DCOVA

- A sample of 11 circuits from a large normal population has a mean resistance of 2.20 ohms. We know from past testing that the population standard deviation is 0.35 ohms.
- Determine a 95% confidence interval for the true mean resistance of the population.



Example

DCOVA


(continued)

- A sample of 11 circuits from a large normal population has a mean resistance of 2.20 ohms. We know from past testing that the population standard deviation is 0.35 ohms.

- **Solution:**

$$\begin{aligned}\bar{X} \pm Z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \\ &= 2.20 \pm 1.96 (0.35/\sqrt{11}) \\ &= 2.20 \pm 0.2068\end{aligned}$$


$$1.9932 \leq \mu \leq 2.4068$$



Interpretation


DCOVA

- We are 95% confident that the true mean resistance is between 1.9932 and 2.4068 ohms
- Although the true mean may or may not be in this interval, 95% of intervals formed in this manner will contain the true mean



Confidence Intervals

DCOVA





Do You Ever Truly Know σ ?

- Probably not!
- In virtually all real world business situations, σ is not known.
- If there is a situation where σ is known then μ is also known (since to calculate σ you need to know μ .)
- If you truly know μ there would be no need to gather a sample to estimate it.



Confidence Interval for μ (σ Unknown)

DCOVA 

- If the population standard deviation σ is unknown, we can substitute the sample standard deviation, S
- This introduces extra uncertainty, since S is variable from sample to sample
- So we use the t distribution instead of the normal distribution

Confidence Interval for μ (σ Unknown)

(continued)

DCOVA A

- Assumptions
 - Population standard deviation is unknown
 - Population is normally distributed
 - If population is not normal, use large sample
- Use Student's t Distribution
- Confidence Interval Estimate:

$$\bar{X} \pm t_{\alpha/2} \frac{S}{\sqrt{n}}$$

(where $t_{\alpha/2}$ is the critical value of the t distribution with $n - 1$ degrees of freedom and an area of $\alpha/2$ in each tail)



Student's t Distribution

DCOVA

- The t is a family of distributions
- The $t_{\alpha/2}$ value depends on **degrees of freedom (d.f.)**
 - Number of observations that are free to vary after sample mean has been calculated

$$\text{d.f.} = n - 1$$

Degrees of Freedom (df)

Idea: Number of observations that are free to vary after sample mean has been calculated

Example: Suppose the mean of 3 numbers is 8.0

Let $X_1 = 7$
Let $X_2 = 8$
What is X_3 ?



If the mean of these three values is 8.0,
then X_3 **must be 9**
(i.e., X_3 is not free to vary)


Here, $n = 3$, so degrees of freedom = $n - 1 = 3 - 1 = 2$

(2 values can be any numbers, but the third is not free to vary for a given mean)

Student's t Distribution

DCOVA


Note: $t \rightarrow Z$ as n increases



Student's t Table

Upper Tail Area			
df	.10	.05	.025
1	3.078	6.314	12.706
2	1.886	2.920	4.303
3	1.638	2.353	3.182

Let: $n = 3$
 $df = n - 1 = 2$
 $\alpha = 0.10$
 $\alpha/2 = 0.05$



The body of the table contains t values, not probabilities

Selected t distribution values

DCOVA

With comparison to the Z value

<u>Confidence Level</u>	<u>t (10 d.f.)</u>	<u>t (20 d.f.)</u>	<u>t (30 d.f.)</u>	<u>Z (∞ d.f.)</u>
0.80	1.372	1.325	1.310	1.28
0.90	1.812	1.725	1.697	1.645
0.95	2.228	2.086	2.042	1.96
0.99	3.169	2.845	2.750	2.58

Note: $t \rightarrow Z$ as n increases

Example of t distribution confidence interval

A random sample of $n = 25$ has $\bar{X} = 50$ and $S = 8$. Form a 95% confidence interval for μ

- d.f. = $n - 1 = 24$, so $t_{\alpha/2} = t_{0.025} = 2.0639$

The confidence interval is

$$\bar{X} \pm t_{\alpha/2} \frac{S}{\sqrt{n}} = 50 \pm (2.0639) \frac{8}{\sqrt{25}}$$

$$46.698 \leq \mu \leq 53.302$$


Example of t distribution confidence interval

(continued)

DCOVAA

- Interpreting this interval requires the assumption that the population you are sampling from is approximately a normal distribution (especially since n is only 25).
- This condition can be checked by creating a:
 - Normal probability plot or
 - Boxplot

Confidence Intervals





Confidence Intervals for the Population Proportion, π

DCOVA A

- An interval estimate for the population proportion (π) can be calculated by adding an allowance for uncertainty to the sample proportion (p)

Confidence Intervals for the Population Proportion, π

(continued)

- Recall that the distribution of the sample proportion is approximately normal if the sample size is large, with standard deviation

$$\sigma_p = \sqrt{\frac{\pi(1-\pi)}{n}}$$

- We will estimate this with sample data:

$$\sqrt{\frac{p(1-p)}{n}}$$

Confidence Interval Endpoints

- Upper and lower confidence limits for the population proportion are calculated with the formula

$$p \pm Z_{\alpha/2} \sqrt{\frac{p(1-p)}{n}}$$

- where
 - $Z_{\alpha/2}$ is the standard normal value for the level of confidence desired
 - p is the sample proportion
 - n is the sample size
- Note: must have $np > 5$ and $n(1-p) > 5$

Example

- A random sample of 100 people shows that 25 are left-handed.
- Form a 95% confidence interval for the true proportion of left-handers



Example

DCOVA

(continued)

- A random sample of 100 people shows that 25 are left-handed. Form a 95% confidence interval for the true proportion of left-handers.

$$\begin{aligned} p \pm Z_{\alpha/2} \sqrt{p(1-p)/n} \\ = 25/100 \pm 1.96 \sqrt{0.25(0.75)/100} \\ = 0.25 \pm 1.96(0.0433) \\ = 0.1651 \leq \pi \leq 0.3349 \end{aligned}$$




Interpretation

- We are 95% confident that the true percentage of left-handers in the population is between
16.51% and 33.49%.
- Although the interval from 0.1651 to 0.3349 may or may not contain the true proportion, 95% of intervals formed from samples of size 100 in this manner will contain the true proportion.



Determining Sample Size

DCOVA





Sampling Error

- The required sample size can be found to reach a desired **margin of error (e)** with a specified level of confidence $(1 - \alpha)$
- The margin of error is also called **sampling error**
 - the amount of imprecision in the estimate of the population parameter
 - the amount added and subtracted to the point estimate to form the confidence interval

Determining Sample Size

Determining
Sample Size

For the
Mean


Sampling error
(margin of error)

$$\bar{X} \pm Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

$$e = Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Determining Sample Size

DCOVA
(continued)



$$e = Z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Now solve
for n to get

$$n = \frac{Z_{\alpha/2}^2 \sigma^2}{e^2}$$

Determining Sample Size

DCOVA
(continued)

- To determine the required sample size for the mean, you must know:
 - The desired level of confidence $(1 - \alpha)$, which determines the critical value, $Z_{\alpha/2}$
 - The acceptable sampling error, e
 - The standard deviation, σ

Required Sample Size Example

DCOVA

If $\sigma = 45$, what sample size is needed to estimate the mean within ± 5 with 90% confidence?

$$n = \frac{Z^2 \sigma^2}{e^2} = \frac{(1.645)^2 (45)^2}{5^2} = 219.19$$

So the required sample size is **$n = 220$**

(Always round up)



If σ is unknown

- If unknown, σ can be estimated when using the required sample size formula
 - Use a value for σ that is expected to be at least as large as the true σ
 - Select a pilot sample and estimate σ with the sample standard deviation, S

Determining Sample Size

(continued)

DCOVA

**Determining
Sample Size**

**For the
Proportion**

$$e = Z \sqrt{\frac{\pi(1-\pi)}{n}}$$

Now solve
for n to get

$$n = \frac{Z_{\alpha/2}^2 \pi (1-\pi)}{e^2}$$



Determining Sample Size

(continued)

DCOVA

- To determine the required sample size for the proportion, you must know:
 - The desired level of confidence ($1 - \alpha$), which determines the critical value, $Z_{\alpha/2}$
 - The acceptable sampling error, e
 - The true proportion of events of interest, π
 - π can be estimated with a pilot sample if necessary (or conservatively use 0.5 as an estimate of π)



Required Sample Size Example

DCOVA

How large a sample would be necessary to estimate the true proportion defective in a large population within $\pm 3\%$, with 95% confidence?

(Assume a pilot sample yields $p = 0.12$)

Required Sample Size Example

(continued)

Solution:

DCOVA

For 95% confidence, use $Z_{\alpha/2} = 1.96$

$e = 0.03$

$p = 0.12$, so use this to estimate π

$$n = \frac{Z_{\alpha/2}^2 \pi (1 - \pi)}{e^2} = \frac{(1.96)^2 (0.12)(1 - 0.12)}{(0.03)^2} = 450.74$$

So use $n = 451$



Ethical Issues

- A confidence interval estimate (reflecting sampling error) should always be included when reporting a point estimate
- The level of confidence should always be reported
- The sample size should be reported
- An interpretation of the confidence interval estimate should also be provided



Confidence Interval Estimation Is Used Frequently In Auditing

- This is an on-line topic




Chapter Summary

In this chapter we discussed

- The concept of confidence intervals
- Point estimates & confidence interval estimates
- Finding confidence interval estimates for the mean (σ known)
- Finding confidence interval estimates for the mean (σ unknown)
- Finding confidence interval estimates for the proportion
- Determining required sample size for mean and proportion
- Ethical issues in confidence interval estimation

*Statistics for Managers Using
Microsoft Excel*
7th Edition



On Line Topic

**Confidence Interval Estimation
In Auditing**



Applications in Auditing

DCOVA

- Six advantages of statistical sampling in auditing
 - Sampling is less time consuming and less costly
 - Sampling provides an objective way to calculate the sample size in advance
 - Sampling provides results that are objective and defensible.
 - Because the sample size is based on demonstrable statistical principles, the audit is defensible before one's superiors and in a court of law.



Learning Objectives

In this section, you learn:

- How to use confidence interval estimates in auditing

Applications in Auditing

DCOVA

(continued)

- Sampling provides an estimate of the sampling error
 - Allows auditors to generalize their findings to the population with a known sampling error.
 - Can provide more accurate conclusions about the population
- Sampling is often more accurate for drawing conclusions about large populations.
 - Examining every item in a large population is subject to significant non-sampling error
- Sampling allows auditors to combine, and then evaluate collectively, samples collected by different individuals.

Confidence Interval for Population Total Amount

- Point estimate for a population of size N:

$$\text{Population total} = N\bar{X}$$

- Confidence interval estimate:

$$N\bar{X} \pm N(t_{\alpha/2}) \frac{S}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}}$$

(This is sampling without replacement, so use the finite population correction in the confidence interval formula)



Confidence Interval for Population Total: Example

DCOVA A

A firm has a population of 1000 accounts and wishes to estimate the total population value.

A sample of 80 accounts is selected with average balance of \$87.6 and standard deviation of \$22.3.

Find the 95% confidence interval estimate of the total balance.

Example Solution

$$N = 1000, \quad n = 80, \quad \bar{X} = 87.6, \quad S = 22.3$$

$$\begin{aligned} N\bar{X} \pm N(t_{\alpha/2}) \frac{S}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}} \\ = (1000)(87.6) \pm (1000)(1.9905) \frac{22.3}{\sqrt{80}} \sqrt{\frac{1000-80}{1000-1}} \\ = 87,600 \pm 4,762.48 \end{aligned}$$

The 95% confidence interval for the population total balance is \$82,837.52 to \$92,362.48

Confidence Interval for Total Difference

- Point estimate for a population of size N:

$$\text{Total Difference} = N\bar{D}$$

- Where the average difference, \bar{D} , is:

$$\bar{D} = \frac{\sum_{i=1}^n D_i}{n}$$

where D_i = audited value - original value

Confidence Interval for Total Difference

(continued)

- Confidence interval estimate:

DCOVA A

$$N\bar{D} \pm N(t_{\alpha/2}) \frac{S_D}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}}$$

where

$$S_D = \sqrt{\frac{\sum_{i=1}^n (D_i - \bar{D})^2}{n-1}}$$

One-Sided Confidence Intervals

- Application: find the **upper bound** for the proportion of items that do not conform with internal controls

$$\text{Upper bound} = p + Z_{\alpha} \sqrt{\frac{p(1-p)}{n}} \sqrt{\frac{N-n}{N-1}}$$

- where
 - Z_{α} is the standard normal value for the level of confidence desired
 - p is the sample proportion of items that do not conform
 - n is the sample size
 - N is the population size




Topic Summary

In this topic we discussed

- Applications of confidence interval estimation in auditing
 - Confidence interval estimation for population total
 - Confidence interval estimation for total difference in the population
 - One-sided confidence intervals for the proportion nonconforming

*Statistics for Managers Using
Microsoft Excel*
7th Edition



On Line Topic

**Estimation & Sample Size
Determination For Finite
Populations**



Topic Learning Objectives

In this topic, you learn:

- When to use a finite population correction factor in calculating a confidence interval for either μ or π
- How to use a finite population correction factor in calculating a confidence interval for either μ or π
- How to use a finite population correction factor in calculating a sample size for a confidence interval for either μ or π

Use A fpc When Sampling More Than 5% Of The Population ($n/N > 0.05$)

DCOVA

Confidence Interval For μ with a fpc

$$\bar{X} \pm t_{\alpha/2} \frac{S}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}}$$

Confidence Interval For π with a fpc

$$p \pm z_{\alpha/2} \sqrt{\frac{p(1-p)}{n}} \sqrt{\frac{N-n}{N-1}}$$

A fpc simply reduces the standard error of either the sample mean or the sample proportion

Confidence Interval for μ with a fpc

DCOVA

Suppose $N = 1000$, $n = 100$, $\bar{X} = 50$, $s = 10$

$$\begin{aligned} 95\% \text{ CI for } \mu &: \bar{X} \pm t_{\alpha/2} \frac{s}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}} = \\ 50 \pm 1.984 \frac{10}{\sqrt{100}} \sqrt{\frac{1000-100}{1000-1}} &= \\ 50 \pm 1.88 &= (48.12, 51.88) \end{aligned}$$

Determining Sample Size with a fpc

DCOVA

- Calculate the sample size (n_0) without a fpc

- For μ :
$$n_0 = \frac{Z_{\alpha/2}^2 \sigma^2}{e^2}$$

- For π :
$$n_0 = \frac{Z_{\alpha/2}^2 \pi(1-\pi)}{e^2}$$


- Apply the fpc utilizing the following formula to arrive at the final sample size (n).
 - $n = n_0 N / (n_0 + (N-1))$



Topic Summary

In this topic we discussed

- When to use a finite population correction in calculating a confidence interval for either μ or π
- The formulas for calculating a confidence interval for either μ or π utilizing a finite population correction
- The formulas for calculating a sample size in a confidence interval for either μ or π



This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher.

Printed in the United States of America.